

Iterative Approaches to Row Sampling ^{*}

Gary L. Miller
CMU

glmiller@cs.cmu.edu

Richard Peng [†]
CMU

yangp@cs.cmu.edu

November 13, 2012

Abstract

We show faster algorithms for solving regression problems based on estimating statistical leverage scores. A growing number of applications involve large, sparse, $n \times d$ matrices $\mathbf{A}$ where $n \gg d$. For many of these applications the more expensive operations involve the $d \times d$ matrix $\mathbf{A}^T \mathbf{A}$. When n is much larger than d , the running time bottleneck is in the cost of computing the matrix $\mathbf{A}^T \mathbf{A}$, rather than the more expensive operations on this smaller matrix.

Recent works by Clarkson, Drineas, Magdon-Ismail, Mahoney and Woodruff led to algorithms that approximate $\mathbf{A}^T \mathbf{A}$ in $O(nd \log n + \text{poly}(d))$ and $O(\text{nnz}(\mathbf{A}) + \text{poly}(d))$ time respectively. When the size of $\mathbf{A}$ is much larger than the cost of more expensive operations involving the smaller $d \times d$ matrix, these algorithms offer significant savings in running time.

We give alternate approaches for approximating $\mathbf{A}^T \mathbf{A}$ based on techniques originally developed for spectral sparsification. Our algorithm finds a matrix $\mathbf{B}$ consisting of a small number of scaled rows of $\mathbf{A}$ such that with high probability $\|\mathbf{A}\mathbf{x}\|_2 = (1 \pm \epsilon)\|\mathbf{B}\mathbf{x}\|_2$ for all vectors $\mathbf{x} \in \mathbb{R}^d$. The running time of our algorithm can be bounded by $O(\text{nnz}(\mathbf{A}) + d^{\omega+\alpha}\epsilon^{-2})$ for any constant $\alpha > 0$. The key to our approach is to find a sequence of estimates of $\mathbf{A}^T \mathbf{A}$ and gradually improve the approximations.

1 Introduction

Least squares calculations and singular vector/value computations are among the most common computational operations involving matrices. These routines are used as core steps in clustering, learning, and more recently algorithmic graph theory. For many operations involving the matrix $\mathbf{A}$, the computationally more expensive steps are often performed on its product, $\mathbf{A}^T \mathbf{A}$. One of the simplest settings where this occurs is with over-determined systems of linear equations of the form $\mathbf{A}\mathbf{x} = \mathbf{b}$. Multiplying both sides by $\mathbf{A}^T$ gives $\mathbf{A}^T \mathbf{A}\mathbf{x} = \mathbf{A}^T \mathbf{b}$, and when $\mathbf{A}$ has full column rank, multiplying both sides by $(\mathbf{A}^T \mathbf{A})^{-1}$ gives $\mathbf{x} = (\mathbf{A}^T \mathbf{A})^{-1} \mathbf{A}^T \mathbf{b}$. This can be shown that this choice of $\mathbf{x}$ minimizes $\|\mathbf{A}\mathbf{x} - \mathbf{b}\|_2$, a common objective in least squares regression [Str93].

Many of the operations involving $\mathbf{A}^T \mathbf{A}$, such as inverse or eigenvector computation, take $O(d^\omega)$ time where ω is the matrix-multiplication constant. When the number of rows, n , is comparable to the number of columns, d , $\mathbf{A}^T \mathbf{A}$ can be computed at a fraction of the cost of these more expensive operations. Also, for iterative methods, the linear operator corresponding to $\mathbf{A}^T \mathbf{A}$ can be implicitly applied in $O(\text{nnz}(\mathbf{A}))$ time via two matrix-vector multiplies involving $\mathbf{A}$ and $\mathbf{A}^T$, where $\text{nnz}(\mathbf{A})$ is the number of non-zero entries. As a result, having access to the linear operator corresponding to $\mathbf{A}^T \mathbf{A}$ is taken for granted in most algorithms. However, many applications involve matrices that are tall and skinny, where n is much larger than d . Here, the d^2 entries in $\mathbf{A}^T \mathbf{A}$ is smaller than the original number of non-zero entries in $\mathbf{A}$ itself. Once n is more than d^2 , the cost of computing $\mathbf{A}^T \mathbf{A}$ is the main running time bottleneck both in

^{*}Partially supported by the National Science Foundation under grant number CCF-1018463.

[†]Supported by a Microsoft Fellowship

theory and in practice. Direct evaluation of $\mathbf{A}^T \mathbf{A}$ takes $O(\text{nnz}(\mathbf{A})d)$ time when $\mathbf{A}$ is sparse, or $O(nd^{\omega-1})$ time when $\mathbf{A}$ is dense. These instances are sufficiently common that practical speedups for finding QR factorizations of tall and skinny matrices have been studied in the distributed [SLHD10, ACD⁺10] and MapReduce settings [CG11].

A more direct way of obtaining speedups is to find faster algorithms that give approximations to $\mathbf{A}^T \mathbf{A}$. The error of approximation between $\mathbf{B}^T \mathbf{B}$ and $\mathbf{A}^T \mathbf{A}$ can be measured either additively or multiplicatively. A multiplicative guarantee is more appealing for several reasons. It can be viewed as relating $\|\mathbf{A}\mathbf{x}\|_2^2$ to $\|\mathbf{B}\mathbf{x}\|_2^2$ for all vectors $\mathbf{x}$. This means that $\mathbf{B}$ can be directly used in place of $\mathbf{A}$ for the overconstrained system of linear systems mentioned earlier. Furthermore, this guarantee is same as the requirement of using $\mathbf{B}^T \mathbf{B}$ as a preconditioner for $\mathbf{A}^T \mathbf{A}$ in iterative methods (see e.g. [Axe94, Saa96]). This means $\mathbf{B}^T \mathbf{B}$ can be used in place of $\mathbf{A}^T \mathbf{A}$ to obtain higher accuracy solutions with only a constant factor overhead. This connection is crucial in practice, and has been used to give speedups on overconstrained systems over the solver in LAPACK [AMT10].

One of the first studied approaches for approximating $\mathbf{A}^T \mathbf{A}$ is to find a matrix $\mathbf{B}$ consisting of a subset of rescaled rows of $\mathbf{A}$, and then compute $\mathbf{B}^T \mathbf{B}$. If $\mathbf{B}$ has fewer rows, $\mathbf{B}^T \mathbf{B}$ can be evaluated faster, and it's also possible for it to remain close to $\mathbf{A}^T \mathbf{A}$. To our knowledge it was first shown by Drineas, Mahoney, and Muthukrishnan that this type of algorithms can be used to speed up L_2 regression [DMM06], albeit in restricted settings. For general matrices, Magdon-Ismail first showed that $\mathbf{B}$ can be found in time faster than directly computing the product $\mathbf{A}^T \mathbf{A}$ [MI10]. Drineas et al. [DMIMW12] gave a much faster algorithm for row sampling that runs in $O(nd \log n + \text{poly}(d, \epsilon))$ time. Recently, Clarkson and Woodruff [CW12] gave algorithms for computing least squares regression and low rank approximations in input sparsity, or $O(\text{nnz}(\mathbf{A}) + \text{poly}(d, \epsilon))$ time.

These algorithms have in common that they apply a sequences of transforms to $\mathbf{A}$ oblivious to its structure, and arrive directly at the high-accuracy $(1 \pm \epsilon)$ approximations. For non-uniform $\mathbf{A}$ concentrated in a few rows, it's clear that these rows are more important. In fact, non-uniform sampling was one of the earliest studied methods [DMM06], and remains the method of choice if key sampling probabilities are provided [DMIMW12]. This paper gives an adaptive algorithm based on row sampling that runs in $O(\text{nnz}(\mathbf{A}) + d^{\omega+\alpha})$ time. Our techniques are motivated by the connection between row sampling and computing statistical leverage scores, and are motivated by works by Spielman and Srivastava on spectral sparsification of graphs [SS08]. In this setting, $\mathbf{A}$ is the edge-vertex incidence matrix of a graph, and subsequent extensions of their work showed that sampling probabilities can be bounded by stretches w.r.t. a spanning tree [KLP12]. Our approaches generalize this notion to stretch w.r.t general matrices, and we denote these matrices that we use to measure stretch as bases. These generalized stretches can be used as sampling probabilities, and we will show that similar matrices can be used in place of each other as bases. This allows us to compute approximate stretches and row samplings in an iterative manner, giving the following result:

Theorem 1.1 *Given a $n \times d$ matrix $\mathbf{A}$ along with failure probability $\delta = d^{-c}$ and allowed error ϵ . For any constant $\alpha > 0$, we can find in $O(\text{nnz}(\mathbf{A}) + d^{\omega+\alpha}\epsilon^{-2})$ time, with probability at least $1 - \delta$, a matrix $\mathbf{B}$ consisting of $O((d^{1+\alpha} + \frac{\text{nnz}(\mathbf{A})}{d^3})\epsilon^{-2})$ rescaled rows of $\mathbf{A}$ such that $\|\mathbf{A}\mathbf{x}\|_2 = (1 \pm \epsilon)\|\mathbf{B}\mathbf{x}\|_2$ for all vectors $\mathbf{x} \in \mathbb{R}^d$.*

The organization of the paper is as follows. Section 2 describes notations used throughout the paper, reviews known results that describe key sampling probabilities, and defines generalized stretch. These results, which are mostly present in earlier works, give that crude estimates of $\mathbf{A}^T \mathbf{A}$ can lead to better ones. Then we show two iterated algorithms for row sampling in Sections 3 and 4. Section 3 gives an algorithm that iterates on projected versions of $\mathbf{A}$, giving the result as stated in Theorem 1.1. Section 4 describes an alternate algorithm that iterates on an artificially perturbed version of $\mathbf{A}$, giving an algorithm with logarithmic dependency the condition number of $\mathbf{A}^T \mathbf{A}$.

2 Background

In this section we outline key results from previous works on row sampling. These works show that statistical leverage scores are closely associated the probabilities needed for row sampling, and give algorithms that efficiently approximate these values. In addition, we formalize the observation implicit in earlier works that both the sampling and estimation algorithms are very robust. The high error-tolerance of these algorithms makes them ideal as core routines to build iterative algorithms upon. Most of the results mentioned in this section are direct adaptations from earlier works, where they're presented in much more detail. For completeness, we give their proofs in Appendix A.

Most of the presentation will use standard linear algebraic notations. We use $\|\mathbf{x}\|_p$ to denote the L_p norm of a vector. The two values of p that we'll use are $p = 1$ and $p = 2$, which correspond to $\|\mathbf{x}\|_1 = \sum_i |x_i|$ and $\|\mathbf{x}\|_2 = \sqrt{\sum_i x_i^2}$. For two vectors $\mathbf{x}$ and $\mathbf{y}$, $\mathbf{x} \geq \mathbf{y}$ means $\mathbf{x}$ is entry-wise greater or equal to $\mathbf{y}$, aka. $x_i \geq y_i$ for all i . Given a matrix $\mathbf{A}$, we use $\mathbf{A}_{i,:}$, or $\mathbf{a}_i$ to denote the i^{th} row of $\mathbf{A}$. Note that $\mathbf{a}_i$ is a row vector of length d . We will also use its Forbenius norm, $\|\mathbf{A}\|_F$ to denote the 2-norm of all its entries. Formally, $\|\mathbf{A}\|_F^2 = \sqrt{\sum_{i,j} \mathbf{A}_{i,j}^2}$.

The spectral theorem states that any symmetric matrix $\mathbf{C}$ has a full complement of orthonormal eigenvectors. Let them and their eigenvalues be $(\mathbf{u}_1, \lambda_1), (\mathbf{u}_2, \lambda_2), \dots (\mathbf{u}_n, \lambda_n)$. Here $\mathbf{u}_j$ is a length d column vector, λ_j is a scalar and they satisfy $\mathbf{A}^T \mathbf{A} \mathbf{u}_j = \lambda_j \mathbf{u}_j$. Without loss of generality we assume that λ_j s are in increasing order: $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$. Then if λ_k is the first non-zero eigenvalue, we have:

$$\mathbf{C} = \sum_{j=k}^n \lambda_j \mathbf{u}_j \mathbf{u}_j^T$$

We will use $\lambda_{\max}$ to denote the largest eigenvalue, and $\lambda_{\min}$ to denote the smallest **non-zero** eigenvalue. Suppose λ_k is the first non-zero eigenvalue, the pseudoinverse of $\mathbf{C}$, $\mathbf{C}^+$ can be defined as:

$$\mathbf{C}^+ = \sum_{j=k}^n \frac{1}{\lambda_j} \mathbf{u}_j \mathbf{u}_j^T$$

This is a linear operator that preserves the null space of $\mathbf{C}$, while acting as its inverse on the rank space. A matrix $\mathbf{C}$ is positive semi-definite if all its eigenvalues are non-negative, or equivalently $\mathbf{x}^T \mathbf{C} \mathbf{x} \geq 0$ for all vectors $\mathbf{x}$. Similarities between matrices is defined using a partial order on matrices. Given two matrices $\mathbf{A}$ and $\mathbf{B}$, $\mathbf{A} \preceq \mathbf{B}$ denotes that $\mathbf{B} - \mathbf{A}$ is positive semi-definite.

We will use $\mathbf{K}$ to specifically to denote $\mathbf{A}^T \mathbf{A}$. An alternate view of row sampling is that we're given n PSD matrices $\mathbf{K}_1 \dots \mathbf{K}_n$ where $\mathbf{K}_i = \mathbf{a}_i^T \mathbf{a}_i$; and the goal is to find a small subset of them such that their weighted sum is close to $\mathbf{K}$. Furthermore, we can assign weights of zero to elements that are not selected, giving a weight vector $\mathbf{w} \in \mathbb{R}^n$. The formal requirement of row sampling can be described as:

- For some parameter κ we have $\mathbf{K} \preceq \sum_{i=1}^n w_i \mathbf{K}_i \preceq \kappa \mathbf{K}$
- $|\{i : w_i \neq 0\}|$, the support of $\mathbf{w}$, is small.

The work by Clarkson and Woodruff [CW12] obtained approximations to $\mathbf{A}$ by applying a (randomized) linear transformation $\mathbf{S}$ to get $\mathbf{B} = \mathbf{S} \mathbf{A}$. In our case, such a matrix can be constructed by letting each row correspond to a non-zero weight in $\mathbf{w}$. This row in $\mathbf{S}$ then has one non-zero entry equaling to the square root of the weight. This is similar to the method used by Clarkson and Woodruff in that each column has at most one non-zero entry, but has the additional property that each row has at most one non-zero entry, that's also non-negative.

2.1 Row Sampling Using Statistical Leverage Scores

Matrix concentration bounds can be viewed as generalizations of single-variable concentration bounds. To our knowledge, the first nearly-tight bounds that encapsulates row sampling was given by Ahlswede and Winter [AW02]. Similar bounds were given as part of a much larger set of results by Rudelson and Vershynin [RV07], although simplifications of their proofs are similar to previous works [Ver09, Har11]. These results identified statistical leverage scores as the crucial numbers corresponding to sampling probabilities. Formally, the statistical leverage score of a row of $\mathbf{A}$, $\mathbf{a}_i$ is defined as:

$$\tau_i = \mathbf{a}_i \mathbf{K}^+ \mathbf{a}_i^T$$

Where $\mathbf{K}^+$ is the pseudoinverse of $\mathbf{K}$.

Although these values have been studied in statistics, their use in algorithms is more recent. Spielman and Srivastava used these bounds in their spectral sparsification algorithm [SS08]. However, their algorithm was restricted to the setting where $\mathbf{A}$ is the edge-vertex incidence matrix of some graph. Works by Drineas, Mahoney, Avron and Toledo generalized this sampling scheme to arbitrary PSD matrices [DM10, AT11]. For our algorithms, we have crude approximations of leverage scores, but only need to bound the number of samples. Such extensions are immediate from the presentations given in [Ver09, Har11]. We use the statement obtained by combining Theorems 5.4. and 6.2. from [AT11], as it gives guarantees for sampling with upper bounds of leverage scores.

Lemma 2.1 (Theorem 5.4, 6.2 from [AT11]) *Let $\mathbf{A}$ be a $n \times d$ matrix and $\mathbf{K}$ be its inner product, $\mathbf{K} = \mathbf{A}^T \mathbf{A}$. Suppose $\tilde{\tau}$ is a length n vector such that $\tilde{\tau}_i \geq \tau_i$. Associate the probability with every row:*

$$p_i = \frac{\tilde{\tau}_i}{\|\tilde{\tau}\|_1} \quad (2.1)$$

and let $\mathbf{T}_1 \dots \mathbf{T}_M$ be i.i.d. random matrices defined by:

$$\mathbf{T}_i = p_{J_i}^{-1} \mathbf{K}_{J_i} \quad (2.2)$$

Where $J_1 \dots J_M$ are random integers between 1 and n which take value i with probability p_i . Then there exist an absolute constant C such that given any error parameter ϵ , choosing $M = C \|\tilde{\tau}\|_1 \log(\|\tilde{\tau}\|_1) / \epsilon^2$ gives:

$$\Pr \left((1 - \epsilon)K \preceq \frac{1}{M} \sum_i \mathbf{T}_i \preceq (1 + \epsilon)K \right) \geq 1 - \frac{1}{\text{poly}(M)}$$

Furthermore, this process, which we denote as $\text{SAMPLE}(\mathbf{A}, \tilde{\tau}, \epsilon)$ can run in $O(n + M \log n)$ time.

Note that the total number of sample depends on the sum of the upper bounds. In the case we use the exact scores, the following fact implies about $O(d \log d)$ samples suffice.

Fact 2.2 (see e.g. [SS08])

$$\sum_{i=1}^n \tau_i = r \leq d$$

The sampling process described in Lemma 2.1 is with replacement, and is similar to the framework described in [AW02, RV07, Har11]. The running time can be obtained by computing partial sums of $\tilde{\tau}$ and binary search for the entry where the cumulative probability meets our threshold. Ipsen and Wentworth

showed that a variety of different sampling methods lead to similar results both theoretically and experimentally [IW12]. Since $\tilde{\tau}_i$ only needs to be upper bounds for τ_i , we can also use poor approximations of leverage scores by scaling them up by the approximation factor. This scaling leads to an increase in the number of rows sampled by the same factor as the approximation ratio. However, it allows us to obtain a much more accurate estimate of $\mathbf{A}$ compared to the accuracy of the approximate statistical leverage scores.

2.2 Generalized Stretch and Their Properties

As our algorithm is iterative and its output is approximate, we will not be able to compute $\mathbf{K}$ exactly. Therefore, all the statistical leverage score estimations in our algorithms will be done using an approximation $\tilde{\mathbf{K}}$ in place of $\mathbf{K}$. As a result, we will incorporate the matrix that we compute these scores by, $\mathbf{K}$ as part of our notation. This is a common notion used in the design of combinatorial preconditioners, where $\tilde{\mathbf{K}}$ is often chosen to be a simpler matrix than $\mathbf{K}$. In the tree setting, Spielman and Woo [SW09] showed that when $\mathbf{A}$ is an edge-vertex incidence matrix and $\tilde{\mathbf{K}}$ corresponds to a spanning tree, $\mathbf{a}_i \tilde{\mathbf{K}} \mathbf{a}_i^T$ equals to the combinatorial stretch of the edge i w.r.t. the tree. This motivates us to use generalized stretch to describe this term for arbitrary matrices. Suppose we're given a factorization of $\tilde{\mathbf{K}}$ as $\mathbf{B}^T \mathbf{B}$, we will use $\mathcal{STR}_{\mathbf{B}}(\mathbf{a}_i)$ to denote:

$$\mathcal{STR}_{\mathbf{B}}(\mathbf{a}_i) := \mathbf{a}_i \tilde{\mathbf{K}}^+ \mathbf{a}_i = \mathbf{a}_i (\mathbf{B}^T \mathbf{B})^+ \mathbf{a}_i \quad (2.3)$$

Note that under this notation, the original definition of statistical leverage score of row i , τ_i equals to $\mathcal{STR}_{\mathbf{A}}(\mathbf{a}_i)$. We will also use this generalized notion to denote leverage scores from this point and on, but still use $\tilde{\tau}$ to denote the approximate bounds of leverage scores computed by our algorithms.

Instead of measuring the stretch of a single row of $\mathbf{A}$, we can also incorporate the stretch of a subset of the rows into our notation. If $\mathbf{A}$ is a matrix with n rows, we use the notation $\mathcal{STR}_{\mathbf{B}}(\mathbf{A})$ to denote:

$$\mathcal{STR}_{\mathbf{B}}(\mathbf{A}) = \sum_{i=1}^n \mathcal{STR}_{\mathbf{B}}(\mathbf{a}_i) \quad (2.4)$$

The following fact is also crucial for our analysis:

Fact 2.3 *Let $\tilde{\mathbf{K}} = \mathbf{B}^T \mathbf{B}$. The generalized stretch of the i^{th} row w.r.t $\mathbf{B}$ equals to its L_2^2 norm under the transformation $\tilde{\mathbf{K}}^{+1/2}$:*

$$\mathcal{STR}_{\mathbf{B}}(\mathbf{a}_i) = \|\tilde{\mathbf{K}}^{+1/2} \mathbf{a}_i^T\|_2^2 = \|\mathbf{a}_i \tilde{\mathbf{K}}^{+1/2}\|_2^2$$

and the total stretch of all rows is

$$\mathcal{STR}_{\mathbf{B}}(\mathbf{A}) = \|\tilde{\mathbf{K}}^{+1/2} \mathbf{A}^T\|_F^2 = \|\mathbf{B} \tilde{\mathbf{K}}^{+1/2}\|_F^2$$

Where $\tilde{\mathbf{K}}^{+1/2}$ is the $1/2$ power of the pseudo-inverse of $\tilde{\mathbf{K}}$ and $\|\cdot\|_F$ denotes the Frobenius norm.

Computing stretch using a different matrix, $\mathcal{STR}_{\mathbf{B}}(\mathbf{A})$, is a standard step when analyzing graph based preconditioners, and done implicitly in [KLP12]. It was shown by Avron, Maymounkov and Toledo that similar matrices can be used in place of each other in least squares regression [AMT10]. The same also holds for generalized stretch, where similar matrices can be used to approximate stretch up to the same factor as their difference. This allows us to define the notion of S -stretch base.

Definition 2.4 *A matrix $\mathbf{B}$ is a S -stretch base for $\mathbf{A}$ if:*

1. $\mathcal{STR}_{\mathbf{B}}(\mathbf{a}_i) \geq \tau_i$

2. $\mathcal{STR}_B(\mathbf{A}) \leq S$

This definition is similar to the objective of the column subset selection problem studied by Avron and Boutsidis [AB12]. Its two conditions are consequences of the requirements on sampling probabilities described in Lemma 2.1. Condition 1 implies that $\mathcal{STR}_A(\mathbf{a}_i)$ can be used as sampling probabilities; while condition 2 means if $\kappa = \text{poly}(d)$, the total number of rows needed for a constant error can be bounded by $O(\kappa d \log d)$. Works on spectral sparsifiers of graphs showed that matrices similar to $\mathbf{A}$, $\mathbf{B}$, can still lead to sampling probabilities [SS08, KLP12]. One sufficient condition is that $\mathbf{B}^T \mathbf{B}$ is spectrally close to $\mathbf{A}^T \mathbf{A}$. It can be formalized using Definition 2.4 as follows:

Lemma 2.5 *For fixed $\mathbf{A}$ and $\mathbf{K} = \mathbf{A}^T \mathbf{A}$, if $\tilde{\mathbf{K}} = \mathbf{B}^T \mathbf{B}$ satisfies:*

$$\frac{1}{\kappa} \mathbf{K} \preceq \tilde{\mathbf{K}} \preceq \mathbf{K}$$

Then $\mathbf{B}$ is a κ -stretch base for $\mathbf{A}$.

Using different bases play crucial roles in both of our algorithms described in Sections 4 and 3.

2.3 Estimation of Generalized Stretch

Fact 2.3 shows that gives that $\mathcal{STR}_A(\mathbf{a}_i)$ is the L_2 norm of a vector. Combining this norm using the Johnson-Lindenstrauss transform (e.g. [Ach01]) the leads to faster approximations of all leverage scores. This was first used by Spielman and Srivastava to approximate effective resistances [SS08]. Direct applications of this reduction, or similar algorithms given in [DMIMW12] lead to a running time of at least $\Omega(\text{nnz}(\mathbf{A}) \log n)$. However, the definition of generalized stretch are only one sided bounds pointwise, which means our estimates can also have large, one sided error. We can use fewer vectors in the projection, and scale up the results to correct potential underestimates. This trades off the coefficient on the leading term at the cost of more elements sampled. This way of remedying one-sided error in the Johnson-Lindenstrauss transform was first shown in [KLP12] and relies on a version of the projection theorem with fewer vectors. We use the following version given as Lemma 7 in [IM98]:

Lemma 2.6 (Lemma 7 from [IM98]) *Let u be a unit vector in $\mathbb{R}^\nu$. For any given positive integers k , let $U_1, \dots, U_k$ be random vectors chosen independently from the ν -dimensional Gaussian distribution $N^\nu(0, 1)$. For $X_i = u^T U_i$, define $W = W(u) = (X_1, \dots, X_k)$ and $L = L(u) = \|W\|_2^2$. Then for any $\beta > 1$:*

1. $E(L) = k$
2. $\Pr(L \geq \beta k) < O(k) \exp(-\frac{k}{2}(\beta - (1 + \ln \beta)))$
3. $\Pr(L \leq k/\beta) < O(k) \exp(-\frac{k}{2}(\beta^{-1} - (1 - \ln \beta)))$

The other observation needed to apply this lemma is that $\tilde{\mathbf{K}}^{+1/2}$ from Fact 2.3 can be replaced by any matrix whose product with its transpose equals to $\tilde{\mathbf{K}}$. As $\tilde{\mathbf{K}} = \mathbf{B}^T \mathbf{B}$, $\tilde{\mathbf{K}}^+ = \tilde{\mathbf{K}}^+ \mathbf{B}^T \mathbf{B} \tilde{\mathbf{K}}^+ = (\mathbf{B} \tilde{\mathbf{K}}^+)^T \mathbf{B} \tilde{\mathbf{K}}^+$. Using this matrix in place of $\tilde{\mathbf{K}}^{+1/2}$ avoids computing the $1/2$ power of $\tilde{\mathbf{K}}$. Pseudocode of our estimation algorithm is shown in Algorithm 1, while the error analysis is nearly identical to the ones given in Section 4 of [SS08] and Section 3.2. of [DMIMW12].

Lemma 2.7 *For each i , with probability at least $1 - \delta$ we have:*

$$\|U \tilde{\mathbf{A}} \tilde{\mathbf{K}}^+ \mathbf{a}_i^T\|_2^2 \geq \frac{1}{\beta} \mathcal{STR}_{\tilde{\mathbf{K}}}(\mathbf{a}_i)$$

Algorithm 1 Algorithm for Upper Bounding Stretch

APPROXSTR($\mathbf{A}, \mathbf{B}, \beta, \delta$)

Input: Matrix $\mathbf{A}$, approximation $\mathbf{B}$ such that $\mathbf{B}^T \mathbf{B}$ is a κ -stretch base for $\mathbf{A}$, value κ , parameter β indicating allowed estimation error. Failure probability δ .

Output: Upper bounds for stretches of rows of $\mathbf{A}$ measured w.r.t. $\tilde{\mathbf{K}}$: $\tilde{\tau}_1 \dots \tilde{\tau}_n$.

 Compute $\tilde{\mathbf{K}} = \mathbf{B}^T \mathbf{B}$

 Compute $\mathbf{K}^+$

 Let $k = O(\log_\beta(1/\delta))$

 Let $\mathbf{U}$ be a $k \times n$ matrix with each entry is picked independently from $N(0, 1)$

 Let $\tilde{\tau}_i$ be $\frac{\beta}{k} \|\mathbf{U} \tilde{\mathbf{A}} \tilde{\mathbf{K}}^+ \mathbf{a}_i^T\|_2^2$.

return $\tilde{\tau}$

Proof We can rewrite $\tilde{\mathbf{K}}^+$ as $\tilde{\mathbf{K}}^+ \tilde{\mathbf{A}}^T \tilde{\mathbf{A}} \tilde{\mathbf{K}}^+$ to get:

$$\begin{aligned}
\mathcal{STR}_{\tilde{\mathbf{K}}}(\mathbf{a}_i) &= \mathbf{a}_i \tilde{\mathbf{K}}^+ \mathbf{a}_i^T \\
&= \mathbf{a}_i \tilde{\mathbf{K}}^+ \tilde{\mathbf{K}} \tilde{\mathbf{K}}^+ \mathbf{a}_i^T \\
&= \mathbf{a}_i \tilde{\mathbf{K}}^+ (\tilde{\mathbf{A}}^T \tilde{\mathbf{A}}) \tilde{\mathbf{K}}^+ \mathbf{a}_i^T \\
&= \mathbf{a}_i \tilde{\mathbf{K}}^+ \tilde{\mathbf{A}}^T \tilde{\mathbf{A}} \tilde{\mathbf{K}}^+ \mathbf{a}_i^T \\
&= \|\tilde{\mathbf{A}} \tilde{\mathbf{K}}^+ \mathbf{a}_i^T\|_2^2
\end{aligned} \tag{2.5}$$

By Lemma 2.6 Part 3 we have:

$$\begin{aligned}
&\Pr \left[\frac{1}{k} \|\mathbf{U} \tilde{\mathbf{A}} \tilde{\mathbf{K}}^+ \mathbf{a}_i^T\|_2^2 < \frac{1}{\beta} \|\tilde{\mathbf{A}} \tilde{\mathbf{K}}^+ \mathbf{a}_i^T\|_2^2 \right] \\
&\leq \exp \left(-\frac{k}{2} (\beta^{-1} - (1 - \ln \beta)) \right) \\
&\leq \exp \left(-\frac{k}{2} (\ln \beta - 1) \right) \\
&\leq \exp \left(-\frac{k \ln \beta}{2} \right) = \beta^{-\frac{k}{2}}
\end{aligned} \tag{2.6}$$

By a suitable choice of constants in $k = O(\log_\beta(1/\delta))$ this can be made less than δ . ■

Bounding the total of $\tilde{\tau}_i$, as well as the running time gives the following guarantees about APPROXSTR:

Lemma 2.8 Let $\mathbf{A}$ be a $n \times d$ matrix, and $\mathbf{B}$ an approximation with $\tilde{n}$ rows such that $\tilde{\mathbf{K}} = \mathbf{B}^T \mathbf{B}$ is a κ -stretch base for $\mathbf{A}$. Then APPROXSTR($\mathbf{A}, \mathbf{B}, \beta, \delta$) runs in $O((\mathbf{nnz}(\mathbf{A}) + \mathbf{nnz}(\tilde{\mathbf{A}}) + d^2) \log_\beta(1/\delta) + (\tilde{n} + d)d^{\omega-1} \log d \epsilon^{-2})$ time returns upper bounds $\tilde{\tau}_i$ such that:

1. For each i , with probability at least $1 - \delta$ we have $\tilde{\tau}_i \geq \tau_i$
2. $\|\tilde{\tau}\|_1 \leq O(\beta \kappa d \log(1/\delta))$ with probability at least $1 - \delta$.

Proof Lemma 2.7 gives

$$\|\mathbf{U} \tilde{\mathbf{A}} \tilde{\mathbf{K}}^+ \mathbf{a}_i^T\|_2^2 \geq \frac{1}{\beta} \mathcal{STR}_{\tilde{\mathbf{K}}}(\mathbf{a}_i)$$

This and Part 1 of Definition 2.4 means scaling the estimates up by a factor of β suffices to give upper bounds of the leverage scores. The upper bound on $\sum_i \tilde{\tau}_i$ then can be obtained similar to the proof of Lemma 2.7, along with the bound on $\mathcal{STR}_{\tilde{\mathbf{K}}}(\mathbf{mata})$ given in Part 2 of Lemma 2.6.

Note that:

$$\begin{aligned} \sum_i \tilde{\tau}_i &= \frac{\kappa\beta}{k} \|\mathbf{U}\tilde{\mathbf{A}}\tilde{\mathbf{K}}^+ \mathbf{a}_i^T\|_2^2 \\ &= \frac{\kappa\beta}{k} \|\mathbf{U}\tilde{\mathbf{A}}\tilde{\mathbf{K}}^+ \mathbf{A}^T\|_F^2 \end{aligned} \quad (2.7)$$

Let $\mathbf{y}$ be the vector that's the row norms of $\mathbf{B}\tilde{\mathbf{K}}^+ \mathbf{A}$. Note that $\|\mathbf{y}\|_2^2 = \sum_{i=1}^n \|\mathbf{B}\tilde{\mathbf{K}}^+ \mathbf{a}_i^T\|_2^2 = \mathcal{STR}_{\tilde{\mathbf{K}}}(\mathbf{A}) \leq \kappa d$, where the last inequality follows from Part 2 of Definition 2.4. Part 2 of Lemma 2.6 can be used to bound $\frac{1}{k} \|\mathbf{U}\mathbf{y}\|_2^2$. Note that when $t > 10$, $t - 1 - \ln t \geq t/2$. If k is larger than some absolute constant, this gives:

$$\begin{aligned} &\Pr\left(\frac{1}{k} \|\mathbf{U}\mathbf{y}\|_2^2 > \beta \kappa d \geq \log(1/\delta) \|\mathbf{y}\|_2^2\right) \\ &\leq O\left(k \exp\left(-\frac{k \log(1/\delta)}{2}\right)\right) \leq \delta \end{aligned} \quad (2.8)$$

It remains to bound the running time of LEVERAGEUPPER. Finding $\mathbf{K}$ takes $O(\tilde{n}d^{\omega-1})$ time, while inverting it takes an additional d^ω time. Computing $\mathbf{U}\tilde{\mathbf{A}}$ can be done in $O(\mathbf{nnz}(\tilde{\mathbf{A}})k)$ time, and multiplying it on the right by $\mathbf{K}^+$ to obtain $\mathbf{U}\tilde{\mathbf{A}}\tilde{\mathbf{K}}^+$ takes $O(kd^2)$ time. It remains to evaluate $\mathbf{U}\tilde{\mathbf{A}}\tilde{\mathbf{K}}^+ \mathbf{a}_i^T$ for each row i . This can be done by summing $\mathbf{nnz}(\mathbf{a}_i)$ length k vectors, giving a total of $\mathbf{nnz}(\mathbf{A})k$ over all n vectors. Therefore the total cost for computing the estimates is $O(k(\mathbf{nnz}(\mathbf{A}) + \mathbf{nnz}(\tilde{\mathbf{A}}) + d^2) + (\tilde{n} + d)d^{\omega-1})$. ■

3 Iterated Row Sampling

In this section we show an algorithm that computes row samplings by computing upper bounds for the leverage scores of blocks of $\mathbf{A}$. It starts from the observation given in Fact 2.3, namely that the stretch of row i is the L_2 norm of the i^{th} row of a matrix, $\mathbf{Y}$. An immediate extension of this is that the generalized stretch of a block of rows of $\mathbf{A}$ equals to the Forbenius norm of the corresponding rows in $\mathbf{Y}$. The Johnson-Lindestrauss lemma as stated in Lemma 2.6 in turn implies that taking a (random) linear combination of these rows preserves the total Forbenius norm. In other words, when the original matrix is used as the base, the generalized stretch of the smaller set of rows w.r.t. $\mathbf{A}$ is comparable to that of the original block. Somewhat surprisingly, switching the base to the new matrix only introduces only a small error. Therefore, the probabilities required by Lemma 2.1 for computing row samplings can be obtained on this smaller matrix. Once again, we need to approximate the product of this smaller matrix to obtain these estimates. However, the significantly reduced row count means we can iterate on it by computing another projection. In a constant number of iterations the number of rows becomes fewer than $\mathbf{nnz}(\mathbf{A})/d$, at which point we can directly compute the answer. We then show that approximate leverage scores computed on these smaller projections can be prolonged to earlier, larger matrices while still meeting the requirements of Lemma 2.1. This leads to a process that resembles V-cycle algorithms from the multigrid literature [BHM00]. These algorithms create a sequence of coarser versions of the initial matrix, and compute answers by going once downward, and once upward through the entire sequence of problems. A representation of our algorithm is given in Figure 1

3.1 Projections and Prolongations

We first formalize the processes of projecting a matrix to one with fewer rows, and prolonging estimates on the projection back to the original matrix. Our key operation is to combine every R rows into k rows, where




Figure 1: High level structure of algorithm: projections to build a sequence of smaller matrices, then prolongations to convert solutions for smaller matrices to solutions for larger ones.

R and k are parameters that we set to d^α and a constant respectively, . By padding $\mathbf{A}$ with additional rows of zeros, we may assume that the number of rows is divisible by R . We will use $n_b = n/R$ to denote the number of blocks, and use the notation (b) to index into the b^{th} block. Projection and prolongation can be formally defined as:

1. (R, k) -projection: for each block $\mathbf{A}_{(b)}$, pick $\mathbf{U}_{(b)}$ to be a $k \times R$ random Gaussian matrix with entries picked independently from $\mathcal{N}(0, 1)$ and compute $\mathbf{A}(short)_{(b)} = \mathbf{U}_{(b)} \mathbf{A}_{(b)}$. Concatenate the blocks $\mathbf{A}(short)_{(b)}$ together vertically to form $\mathbf{A}(short)$.
2. (k, R) -prolongation: given a vector $\tilde{\tau}'$ with length $n_b k$, maps it to length $n_b R$ vector where each entry in block b has value $\|\tilde{\tau}'_{(b)}\|_1$.

We first show that projections preserve the stretch of blocks w.r.t. $\mathbf{A}$. This can be done by bounding the effect of $\mathbf{U}_{(b)}$ on the norm of each column of $\mathbf{A}_{(b)} \mathbf{K}^{+1/2}$. It follows directly properties of Johnson-Lindenstrauss projections described in Lemma 2.6, and a proof is given in Appendix B.

Lemma 3.1 *Assume $R \geq d^\alpha$ for some constant α and let $\mathbf{A}(short)$ be a (R, k) -projection of $\mathbf{A}$. For any constant $c > 0$ there exists a constant $k = O(c/\alpha)$ such that for any block b , we have with probability at least $1 - d^{-c}$:*

$$\mathcal{STR}_{\mathbf{A}}(\mathbf{A}(short)_{(b)}) \geq \frac{k}{R} \mathcal{STR}_{\mathbf{A}}(\mathbf{A}_{(b)})$$

We next show that we can change the base from $\mathbf{A}$ to $\mathbf{A}(short)$, and use $\mathcal{STR}_{\mathbf{A}(short)}(\mathbf{A}(short)_{(b)})$ as upper bounds for $\mathcal{STR}_{\mathbf{A}}(\mathbf{A}(short)_{(b)})$. As a first step, we need to relate $\mathbf{K}$ to the product of $\mathbf{A}(short)$, $\mathbf{K}(short) = \mathbf{A}(short)^T \mathbf{A}(short)$. Since each $\mathbf{A}(short)_{(b)}$ is formed by merging rows of $\mathbf{A}_{(b)}$, $\mathbf{A}(short)_{(b)} = \mathbf{U}_{(b)} \mathbf{A}_{(b)}$, $\mathbf{A}(short)_{(b)}^T \mathbf{A}(short)_{(b)}$ can be upper bounded by $\|\mathbf{U}_{(b)}\|_F^2 \cdot \mathbf{A}_{(b)}^T \mathbf{A}_{(b)}$.

Lemma 3.2 *The following holds for each block b :*

$$\mathbf{A}(short)_{(b)}^T \mathbf{A}(short)_{(b)} \preceq \|\mathbf{U}_{(b)}\|_F^2 \cdot \mathbf{A}_{(b)}^T \mathbf{A}_{(b)}$$

A proof is given in Appendix B. Generalized stretches w.r.t. $\mathbf{K}(short)$ and $\mathbf{K}$ are evaluated under the norms given by $\mathbf{K}(short)^+$ and $\mathbf{K}^+$. Therefore, we need the inverse of this bound, which we obtain using the following lemma about pseudoinverses.

Lemma 3.3 *Let $\mathbf{C}$ and $\mathbf{D}$ be symmetric positive semi-definite matrices and let Π be the projection space onto the range space of $\mathbf{C}$. Then:*

$$\Pi \mathbf{C}^+ \Pi \succeq \Pi (\mathbf{C} + \mathbf{D})^+ \Pi$$

This statement is straightforward when both $\mathbf{C}$ and $\mathbf{D}$ are full rank, or share the same null space. However, as pseudo-inverses do not act on the null space, it is crucial that we're only considering vectors of the form $\mathbf{a}'_i$. This Lemma is also proven in Appendix B. Combining it with bounds on $\|\mathbf{U}_{(b)}\|_F^2$ allows us to bound the distortion caused by switching the base matrix from $\mathbf{A}$ to $\mathbf{A}(\text{short})$.

Lemma 3.4 *For any constant c , there exists a constant c' such that with probability at most $1 - d^{-c}$, for any row i of $\mathbf{A}(\text{short})$, $\mathbf{a}(\text{short})_i$ we have:*

$$c' \log nkR \cdot \mathcal{STR}_{\mathbf{A}(\text{short})}(\mathbf{a}(\text{short})_i) \geq \mathcal{STR}_{\mathbf{A}(\text{short})}(\mathbf{a}(\text{short})_i)$$

Proof Since each $\mathbf{U}(b)$ consists of $k \times R$ independent random variables chosen from $N(0, 1)$, $\|\mathbf{U}(b)\|_F^2$ is distributed as $N(0, kR)$. This gives:

$$\Pr(\|\mathbf{U}(b)\|_F^2 > l \cdot kR) \leq \exp(-l) \quad (3.9)$$

There exists c' such that setting $l = c' \log n$ allows us to upper bound this probability by n^{-2c} . Taking a union bound over all n/d blocks gives that with probability at least $1 - d^{-c}$ this holds for all blocks.

As before, let $\mathbf{K} = \mathbf{A}^T \mathbf{A}$ and $\mathbf{K}(\text{short}) = \mathbf{A}(\text{short})^T \mathbf{A}(\text{short})$. Applying Lemma 3.2 to all blocks and summing over all blocks gives:

$$\mathbf{K}(\text{short}) \preceq c' \log nkR \cdot \mathbf{K} \quad (3.10)$$

Let Π be the projection operator onto the range space of $\mathbf{K}(\text{short})$, Applying Lemma 3.3 with $\mathbf{C} = \mathbf{K}(\text{short})$ and $\mathbf{D} = c' \log nkR \cdot \mathbf{K}$ gives:

$$\Pi \mathbf{K}(\text{short})^+ \Pi \succeq \frac{1}{c' \log nkR} \Pi \mathbf{K}^+ \Pi \quad (3.11)$$

Note that $\mathbf{a}(\text{short})_i$ is completely contained within the range space of $\mathbf{K}(\text{short})$. Therefore $\Pi \mathbf{a}(\text{short})_i^T = \mathbf{a}(\text{short})_i^T$ and:

$$\begin{aligned} & c' \log nkR \cdot \mathcal{STR}_{\mathbf{A}(\text{short})}(\mathbf{a}(\text{short})_i) \\ &= c' \log nkR \cdot \mathbf{a}(\text{short})_i \mathbf{K}(\text{short})^+ \mathbf{a}_i^T \\ &= c' \log nkR \cdot \mathbf{a}(\text{short})_i \Pi \mathbf{K}(\text{short})^+ \Pi \mathbf{a}(\text{short})_i^T \\ &\geq \mathbf{a}(\text{short})_i \Pi \mathbf{K}^+ \Pi \mathbf{a}(\text{short})_i^T \quad \text{By Lemma 3.3} \\ &= \mathbf{a}(\text{short})_i \mathbf{K}^+ \mathbf{a}(\text{short})_i^T = \mathcal{STR}_{\mathbf{A}}(\mathbf{a}(\text{short})_i) \end{aligned} \quad (3.12)$$

Summing over all k rows of $\mathbf{A}(\text{short})_{(b)}$ completes the proof. $\blacksquare$

Combining Lemmas 3.4 and 3.1 shows that with high probability, scaling up $\mathcal{STR}_{\mathbf{A}(\text{short})}(\mathbf{A}(\text{short})_{(b)})$ by $O(R^2 \log n)$ gives upper bounds for $\mathcal{STR}_{\mathbf{A}}(\mathbf{a}_{(b)})$.

Corollary 3.5 *For any parameter $R \geq d^\alpha$, constant c , there exist constants k and c' such that for a block b , with probability at least $1 - d^{-c}$,*

$$c' R^2 \log n \mathcal{STR}_{\mathbf{A}(\text{short})}(\mathbf{A}(\text{short})_{(b)}) \geq \mathcal{STR}_{\mathbf{A}}(\mathbf{a}_{(b)})$$

3.2 Iterative Algorithm

Projecting $\mathbf{A}$ to $\mathbf{A}(\text{short})$ gives a matrix with fewer rows, and a way to reduce the sizes of our problems. A fast algorithm follows by examining the sequence of matrices $\mathbf{A} = \mathbf{A}(0), \mathbf{A}(1), \dots, \mathbf{A}(L)$ obtained using

such projections. Once $\mathbf{A}(L)$ has fewer than $\mathbf{nnz}(\mathbf{A})d^{-3}$ rows, $\mathbf{A}(L)^T \mathbf{A}(L)$ can be approximated directly. This then allows us to approximate the statistical leverage scores of the rows of $\mathbf{A}(L)$. Corollary 3.5 shows that statistical leverage scores in $\mathbf{A}(l)$, $\tau(l)$ can upper bound those of $\mathbf{A}(l-1)$, $\tau(l-1)$. This means we can gradually prolongate solutions backwards from $\mathbf{A}(L)$ to $\mathbf{A}(0)$. We do so by maintaining the invariant that $\mathbf{B}(l)$ has a small number of rows and is close to $\mathbf{A}(l)$. This means stretches of $\mathbf{A}(l)$ w.r.t. $\mathbf{B}(l)$ as base can be used in place of statistical leverage scores of $\mathbf{A}(l-1)$. These scores can then be used to compute $\mathbf{B}(l-1)$, keeping the invariant for $l-1$. Pseudocode of the algorithm is shown in Algorithm 2.

Algorithm 2 Row Sampling Via. Projections

ROWCOMBINE($\mathbf{A}, R, \epsilon, \delta, n_{\min}$)

Input: Reduction rate R , $n \times d$ matrix $\mathbf{A}$, allowed approximation error ϵ , failure probability $\delta = d^{-c}$.

Output: Sparsifier $\mathbf{B}$ that contains $O(dR^4 \log d / \epsilon^2)$ scaled rows of $\mathbf{A}$ such that $(1 - \epsilon)\mathbf{A}^T \mathbf{A} \preceq \mathbf{B}^T \mathbf{B} \preceq (1 + \epsilon)\mathbf{A}^T \mathbf{A}$.

Set $L = d^3 \log_R$, and fix constants c', k based on c

Set $\epsilon(0) = \epsilon$, $\epsilon(1) \dots \epsilon(l) = 1/2$

$\mathbf{A}(0) = \mathbf{A}$

for $l = 1 \dots L$ **do**

Let $\mathbf{A}(l)$ be a (R, k) -projection of $\mathbf{A}(l-1)$

end for

$\mathbf{B}(L) \leftarrow \mathbf{A}(L)$

for $i = L \dots 1$ **do**

$\tilde{\tau}'(l) \leftarrow \frac{1}{d^3} + O(R^3 \log n) \cdot \text{APPROXSTR}(\mathbf{A}(l), \sqrt{\frac{2}{3}}\mathbf{B}(l), R, d^{-c-6})$

Let $\tilde{\tau}(l-1)$ be a (k, R) -prolongation of $\tilde{\tau}'(l)$

$\mathbf{B}(l-1) \leftarrow \text{SAMPLE}(\mathbf{A}(l-1), \tilde{\tau}(l-1), \epsilon(l))$

end for

return $\mathbf{B}^{(0)}$

The algorithm obtains sampling probabilities for $\mathbf{A}(l-1)$ using (k, R) prolongations of estimates of stretches of $\mathbf{A}(l)$ w.r.t. $\mathbf{B}(l)$. We first show these values, $\tilde{\tau}(l-1)$, are with high probability upper bounds for the statistical leverage scores of $\mathbf{A}(l-1)$, $\tau(l-1)$.

Lemma 3.6 *Suppose $\mathbf{B}(l)$ satisfies:*

$$\frac{1}{2}\mathbf{A}(l)^T \mathbf{A}(l) \preceq \mathbf{B}(l)^T \mathbf{B}(l) \preceq \frac{3}{2}\mathbf{A}(l)^T \mathbf{A}(l)$$

Then with probability at least $1 - 2d^{-c-1}$, we have

- $\tilde{\tau}(l-1) \geq \tau(l-1)$
- $\|\tilde{\tau}(l-1)\|_1 \leq O(dR^4 \log n + \frac{n}{d^3})$

Proof By Lemma 2.5, $\sqrt{\frac{2}{3}}\mathbf{B}(l)$ is a $3d$ -stretch base for $\mathbf{A}(l+1)$. Since $\mathbf{A}(l)$ is a projection of $\mathbf{A}(l-1)$, we can index corresponding blocks in them. Consider any block (b) . Applying Lemma 2.8 to all $R \leq d$ rows of $\mathbf{A}(l)_{(b)}$ gives that we have with probability at least $1 - d^{-c-5}$:

$$R\|\tilde{\tau}'(l)_{(b)}\|_1 \geq \text{STR}_{\mathbf{A}(l)}(\mathbf{A}(l)_{(b)}) \tag{3.13}$$

Combining with Corollary 3.5 applied to $\mathbf{A}(l-1)$ and $\mathbf{A}(l)$ gives that with probability at least $1 - 2d^{-c-5}$ we have:

$$c'R^3 \log n \|\tilde{\tau}'(l)_{(b)}\|_1 \geq \|\tau(l-1)_{(b)}\|_1 \quad (3.14)$$

Also, note that $\frac{1}{d^3}$ is added to all entries of $\tilde{\tau}(l)$. Since $\|\tau(l)\|_1 \leq d$, the total number of blocks where $\frac{1}{d^3}$ does not suffice as an upper bound for some row is at most d^4 . Applying the above bound to these blocks gives that with probability of at least $1 - d^{-c-1}$ we have $\tilde{\tau}(l-1) \geq \tau(l-1)$.

It remains to upper bound $\|\tilde{\tau}(l-1)\|_1$. Lemma 2.8 Part 2 gives with probability at least $1 - d^{-c-6}$, $\|\tilde{\tau}'(l)\|_1 \leq O(d)$. As each $\|\tilde{\tau}'(l)_{(b)}\|_1$ is prolonged to R rows of $\tilde{\tau}(l-1)$, we get:

$$\begin{aligned} \|\tilde{\tau}(l-1)\|_1 &= \sum_b R \left(\frac{1}{d^3} + C \cdot R^3 \log n \|\tilde{\tau}'(l)_{(b)}\|_1 \right) \\ &\leq O \left(\frac{n}{d^3} + R^4 \log n \sum_b \|\tilde{\tau}'(l)_{(b)}\|_1 \right) \\ &\leq O \left(dR^4 \log n + \frac{n}{d^3} \right) \end{aligned} \quad (3.15)$$

The overall probability of $1 - d^{-c-1}$ follows applying union bound once again. $\blacksquare$

Combining these with the fact that the number of rows decrease by a factor of $O(R)$ per iteration completes the algorithm. The bound given in Theorem 1.1 is then obtained by setting R to $d^{\alpha/4}$. Applying 3.6 inductively backwards in l gives the overall bound, and in turn Theorem 1.1.

Lemma 3.7 *If ROWCOMBINE is ran with $R = d^{\alpha/4}$, $\delta = d^{-c}$ where α and c are positive constants, then with probability at least $1 - d^{-c}$ it returns in $O(\mathbf{nnz}(\mathbf{A}) + d^{\omega+\alpha}\epsilon^{-2})$ time $\mathbf{B}$ such that:*

$$(1 - \epsilon)\mathbf{A}^T \mathbf{A} \preceq \mathbf{B}^T \mathbf{B} \preceq (1 + \epsilon)\mathbf{A}^T \mathbf{A}$$

and $\mathbf{B}$ has $\tilde{O}((d^{1+\alpha} \log n + \frac{n}{d^3}) \log d\epsilon^{-2})$ rows, each being a scaled copy of some row of $\mathbf{A}$,

Proof We first show correctness via. induction backwards on l . Specifically we show that with probability at least $1 - 3(L-l)d^{-c-1}$, $\mathbf{B}(l)$ has $O((dR^4 \log n + \frac{n}{d^3}) \log d\epsilon(l)^{-2})$ rows and satisfies $(1 - \epsilon(l))\mathbf{A}(l)^T \mathbf{A}(l)^T \preceq \mathbf{B}(l)^T \mathbf{B}(l) \preceq (1 + \epsilon(l))\mathbf{A}(l)^T \mathbf{A}(l)^T$. Note that since $R = d^\alpha$, $L = \log_R d^3$ is a constant. As the number of rows decreases by a factor of $O(R)$ per projection, we get that $\mathbf{A}(L)$ has $O(n/d^3)$ rows. Therefore the base case where $l = L$ follows from $\mathbf{B}(L) = \mathbf{A}(L)$.

For the inductive step, we assume that the inductive hypothesis holds for $l \geq 1$ and try to show it for $l-1$. As $\epsilon(l)$ was set to $1/2$, we have:

$$\frac{1}{2}\mathbf{A}(l)^T \mathbf{A}(l) \preceq \mathbf{B}(l)^T \mathbf{B}(l) \preceq \frac{3}{2}\mathbf{A}(l)^T \mathbf{A}(l)$$

This allows us to invoke Lemma 3.6, which combined with Lemma 2.1 gives that with probability $1 - d^{-c-1}$, the inductive hypothesis also holds for $l-1$. The failure probability follows from union bounding this with the failure probabilities from the hypothesis and Lemma 3.6.

We now bound the total running time, starting with the projections. Since $\mathbf{U}_{(b)}$ has $k = O(1)$ rows, the total cost of computing $\mathbf{U}_{(b)}\mathbf{A}(l)_{(b)}$ is $O(\mathbf{nnz}(\mathbf{A}(l)))$. Furthermore, we also get $\mathbf{nnz}(\mathbf{A}(l+1)) = O(\mathbf{nnz}(\mathbf{A}(l)))$, giving a total cost of $O(\mathbf{nnz}(\mathbf{A}))$ over all L projections. Since $\mathbf{B}(l)$ has $O((dR^4 \log n + \frac{n}{d^3}) \log d\epsilon(l+1)^{-2})$ rows, Lemma 2.8 gives that each call to APPROXSTR takes $O(d^\omega R^4 \log n \log d\epsilon^{-2} + nd^{\omega-3} \log d\epsilon^{-2})$ time. By standard assumptions that $\log d\epsilon^{-2} \leq d^{3-\omega}$ and $n \leq \mathbf{nnz}(\mathbf{A})$, this is at most $O(d^\omega R^4 \log n \log d\epsilon^{-2} + \mathbf{nnz}(\mathbf{A}))$. Substituting $R = d^{\alpha/4}$ into these bounds gives the result. $\blacksquare$

4 Introduction and Removal of Artificial Bases

This section describes a different algorithm that computes row samplings of $\mathbf{A}$ using artificial bases. These bases are formed by adding d additional rows to $\mathbf{A}$ equaling to $\gamma\mathbf{I}$. When γ very large, this matrix can be shown to be almost the same as the identity matrix. On the other hand, when γ is very small, there is very little difference between this matrix and $\mathbf{A}$. As a result, the key is to show that γ can be reduced geometrically over the iterations. Given an initial value $\gamma(0)$, and a parameter R indicating the reduction rate, we consider a geometrically decreasing sequence of $\gamma(l)$ s defined by $\gamma(l+1) = R^{-1/2}\gamma(l)$. Then $\mathbf{A}(l)$ can be written as:

$$\mathbf{A}(l) = \begin{bmatrix} \mathbf{A} \\ \gamma(l)\mathbf{I} \end{bmatrix}$$

And we will use $\mathbf{K}(l)$ to denote its product:

$$\mathbf{K}(l) = \mathbf{A}(l)^T \mathbf{A}(l) = \mathbf{K} + \gamma(l)^2 \mathbf{I}$$

We will use $\mathbf{B}(l)$ to denote approximations of $\mathbf{A}(l)$ obtained from row samplings, and $\tilde{\mathbf{K}}(l)$ to be the product of $\mathbf{B}(l)$, $\mathbf{B}(l)^T \mathbf{B}(l)$. A geometrically decreasing sequence of values of γ are of special interest to us. We first show that if $\gamma(0)^2$ is more than the maximum eigenvalue of $\mathbf{A}^T \mathbf{A}$, a copy of the identity matrix suffices as $\mathbf{B}(0)$.

Lemma 4.1 *Let $\lambda_{\max}$ be the maximum eigenvalue of $\mathbf{A}^T \mathbf{A}$. If $\gamma(0)^2 \geq \lambda_{\max}$, then $\mathbf{B}(l) = \gamma(0)\mathbf{I}$ satisfies:*

$$\tilde{\mathbf{K}}(0) \preceq \mathbf{K}(0) \preceq 2\tilde{\mathbf{K}}(0)$$

Proof The LHS follows from $\mathbf{K}(l) = \mathbf{K} + \gamma(0)^2 \mathbf{I} \succeq \gamma(0)^2 \mathbf{I} = \mathbf{B}(0)^T \mathbf{B}(0)$. The RHS can be obtained by $\mathbf{K}(0) \succeq \lambda_{\max} \mathbf{I} \succeq \gamma(0)^2 \mathbf{I} \succeq \tilde{\mathbf{K}}(0)$ and adding $\gamma(0)^2 \mathbf{I}$ to both sides. ■

Also, once $\gamma(l)^2$ is less than the minimum eigenvalue of $\mathbf{A}^T \mathbf{A}$, $\mathbf{B}(l)$ can be directly used as a low stretch base for $\mathbf{A}(l)$.

Lemma 4.2 *Let $\lambda_{\min}$ be the minimum eigenvalue of $\mathbf{A}^T \mathbf{A}$. If $\gamma(l)^2 \leq \lambda_{\min}$, then $\sqrt{\frac{1}{2}}\mathbf{B}(l)$ is a $2d$ -stretch base for $\mathbf{A}$.*

Proof For this proof it is easier to work with the product of $\sqrt{\frac{1}{2}}\mathbf{B}(l)$, since $(\sqrt{\frac{1}{2}}\mathbf{B}(l))^T (\sqrt{\frac{1}{2}}\mathbf{B}(l)) = \frac{1}{2}\tilde{\mathbf{K}}(l)$. Let a spectral decomposition of $\mathbf{K}$ be $\mathbf{K} = \sum_{j=1}^d \lambda_j \mathbf{v}_j \mathbf{v}_j^T$, where $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_d$. Assume λ_k is the first non-zero eigenvalue, then:

$$\begin{aligned} \mathbf{K}(l) &= \mathbf{K} + \gamma(l)^2 \mathbf{I} = \sum_{j=k}^d (\lambda_j + \gamma(l)^2) \mathbf{v}_j \mathbf{v}_j^T \\ \frac{1}{2}\mathbf{K}(l)^+ &= \sum_{j=k}^d \frac{1}{\lambda_j + \gamma(l)^2} \mathbf{v}_j \mathbf{v}_j^T \end{aligned}$$

Consider the eigenvector $\mathbf{v}_j$ for some $j < k$, $\lambda_j = 0$ implies $\|\mathbf{A}\mathbf{v}_j\|_2^2 = \mathbf{v}_j^T \mathbf{K} \mathbf{v}_j = 0$, Therefore for any row i of $\mathbf{A}$, we have $\mathbf{a}_i \mathbf{v}_j = 0$. This means for both $\mathbf{a}_i \mathbf{K}^+ \mathbf{a}_i^T$ and $\mathbf{a}_i \mathbf{K}(l)^+ \mathbf{a}_i^T$ it suffices to consider the terms from k to d . For each of these terms, $\gamma \leq \lambda_j$ gives $\frac{1}{2}(\lambda_j + \gamma) \leq \lambda_j$ and therefore:

$$\frac{1}{\lambda_j} \leq \frac{1}{(1/2)(\lambda_j + \gamma)} \quad (4.16)$$

As each term from k to d is non-negative, summing over them meets Part 1 of Definition 2.4.

For the upper bound required by Part 2, note that $\lambda_j \leq \lambda_j + \gamma$ gives:

$$\frac{1}{(1/2)(\lambda_j + \gamma)} \leq \frac{2}{\lambda_j}$$

For a row i of $\mathbf{A}$, $\mathbf{a}_i$, summing over each eigenvalue gives:

$$\mathcal{STR}_{\sqrt{\frac{1}{2}}\mathbf{B}(l)}(\mathbf{a}_i) \leq 2\mathcal{STR}_{\mathbf{A}}(\mathbf{a}_i)$$

. Summing once again over all rows leads to $\mathcal{STR}_{\sqrt{\frac{1}{2}}\mathbf{B}(l)}(\mathbf{A}) \leq 2\mathcal{STR}_{\mathbf{A}}(\mathbf{A}) \leq 2d$. ■

The algorithm is based on the fact that $\mathbf{K}(l+1)$ can be used as $O(dR)$ -stretch base for $\mathbf{K}(l)$. By the requirements of Lemma 2.5, it suffices to show that $\mathbf{K}(l)$ and $\mathbf{K}(l+1)$ are spectrally close.

Lemma 4.3

$$\mathbf{K}(l+1) \preceq \mathbf{K}(l) \preceq R\mathbf{K}(l+1)$$

Proof The LHS then follows from $\mathbf{I}$ being PSD, while for the RHS we have:

$$\begin{aligned} R\mathbf{K}(l+1) &= R(\mathbf{K} + \gamma(l+1)^2\mathbf{I}) \\ &= R(\mathbf{K} + R^{-1}\gamma(l)^2\mathbf{I}) \\ &\succeq \mathbf{K} + \gamma(l)\mathbf{I} \\ &= \mathbf{K}(l) \end{aligned} \tag{4.17}$$

■

Note that if $\frac{1}{2}\mathbf{K}(l) \preceq \tilde{\mathbf{K}}(l) \preceq \frac{3}{2}\mathbf{K}(l)$, Lemma 4.3 implies that

$$\frac{1}{3R}\mathbf{K}(l+1) \preceq \frac{2}{3R}\tilde{\mathbf{K}}(l) \preceq \mathbf{K}(l+1) \tag{4.18}$$

Therefore by Lemma 2.5, $\sqrt{\frac{2}{3R}}\mathbf{B}(l)$ is a $3dR$ -stretch base for $\mathbf{K}(l+1)$. If we're given bounds on non-zero eigenvalues λ_{upper} and λ_{lower} , we can start with $\gamma(0) = \lambda_{upper}$ and continue until $\gamma(l) \leq \lambda_{lower}$. This leads to the algorithm whose pseudocode is given in Algorithm 3

Theorem 4.4 *Let c be any constant corresponding to a failure probability of d^{-c} . Given a $n \times d$ matrix $\mathbf{A}$, upper and lower bounds on the non-zero eigenvalues of $\mathbf{A}^T\mathbf{A}$ λ_{upper} and λ_{lower} , as well as parameters ϵ . ROWSAMPLE runs in $O(((nnz(\mathbf{A}) + d^2) \log_R n + d^\omega R^2 \log^2 d) \log_R(\lambda_{upper}/\lambda_{lower}) + d^\omega R^2 \log^2 d \epsilon^{-2})$ time, and returns a matrix $\mathbf{B}$ that contains $O(dR^2 \log d \epsilon^{-2})$ rescaled rows of $\mathbf{A}$ and $\tilde{\mathbf{K}} = \mathbf{B}^T\mathbf{B}$. With probability at least $1 - d^{-c}$ we have:*

$$(1 - \epsilon)\mathbf{K} \preceq \tilde{\mathbf{K}} \preceq (1 + \epsilon)\mathbf{K}$$

Proof We may assume that $\log_R(\lambda_{upper}/\lambda_{lower}) \leq 1/10d$ since otherwise directly computing $\mathbf{A}^T\mathbf{A}$ gives the required runtime bound.

We first show that $\tilde{\mathbf{K}}(l)$ are good approximations of $\mathbf{K}(l)$ with high probability. We prove by induction on l that with probability $1 - O(ld^{-c-1})$ we have:

$$\frac{1}{2}\mathbf{K}(l) \preceq \tilde{\mathbf{K}}(l) \preceq \frac{3}{2}\mathbf{K}(l) \tag{4.19}$$

Algorithm 3 Row Sampling via. Introduction and Removal of Identity

ROWSAMPLE($R, \lambda_{lower}, \lambda_{upper}, \mathbf{A}, \epsilon, \delta$)

Input: Parameter $R \geq 4$ indicating the speed of reduction. Matrix $\mathbf{A}$, upper and lower bounds on eigenvalues, $\lambda_{upper} \geq \lambda_{\max}(\mathbf{A})$ and $\lambda_{lower} \leq \lambda_{\min}(\mathbf{A})$. Error parameter ϵ and failure probability δ .

Output: Matrix $\mathbf{B}$ containing $O(dR^2 \log d)$ scaled rows of $\mathbf{A}$ such that $(1 - \epsilon)\mathbf{A}^T \mathbf{A} \preceq \mathbf{B}^T \mathbf{B} \preceq (1 + \epsilon)\mathbf{A}^T \mathbf{A}$

```

1: Let  $\gamma(0) = \lambda_{upper}$ 
2: Let  $\mathbf{B}(0) = \sqrt{\lambda_{upper}} \mathbf{I}$ ,  $\mathbf{K}(0) = \lambda_{upper} \mathbf{I}$ ,  $l = 0$ 
3: while  $\gamma(l) > \lambda_{lower}$  do
4:    $l \leftarrow l + 1$ 
5:    $\gamma(l) \leftarrow R^{-1/2} \gamma(l - 1)$ 
6:    $\tilde{\tau}(l) \leftarrow \text{APPROXSTR}(\mathbf{A}(l), \sqrt{\frac{2}{3R}} \mathbf{B}(l - 1), R, d^{-c} n^{-1})$ 
7:    $\mathbf{B}(l) \leftarrow \text{SAMPLE}(\mathbf{A}(l), \tilde{\tau}(l), 1/2, d^{-1} \delta)$ 
8: end while
9:  $\tilde{\tau} \leftarrow \text{APPROXSTR}(\mathbf{A}, \sqrt{\frac{2}{3R}} \mathbf{B}(l), R, d^{-c} n^{-1})$ 
10:  $\mathbf{B} \leftarrow \text{SAMPLE}(\mathbf{A}, \tilde{\tau}, \epsilon, d^{-1} \delta)$ 
11: return  $\mathbf{B}$ 

```

and $\tilde{\mathbf{K}}(l)$ contains $O(dR^2 \log^2 d)$ scaled rows of $\mathbf{A}(l)$. The base case of $l = 0$ follows from Lemma 4.1. For the inductive case, combining Lemma 4.3 and the inductive hypothesis gives:

$$\frac{1}{3R} \mathbf{K}(l) \preceq \frac{2}{3R} \tilde{\mathbf{K}}(l - 1) \preceq \mathbf{K}(l) \quad (4.20)$$

Therefore Lemma 2.5 gives that $\frac{2}{3R} \mathbf{B}(l - 1)$ is an $3dR$ -stretch base for $\mathbf{A}(l)$. Lemma 2.8 then guarantees that for each row i of $\mathbf{A}(l)$, with probability at least $1 - d^{c-1}(n + d)^{-1}$, $\tilde{\tau}(l) \geq \tau(l)$. Applying union bound over all $n + d$ rows of $\mathbf{A}(l)$ gives that with probability at least $1 - d^{-c-1}$ this holds for all rows. Also, with probability at least $1 - d^{-c-1}$, $\|\tilde{\tau}(l)\|_1 \leq O(dR^2 \log(d^{-c-1})) = O(dR^2 \log d)$. If both of these bounds hold, by Lemma 2.1, we have with probability at least $1 - O(d^{-c-1})$, $\frac{1}{2} \mathbf{K}(l) \preceq \tilde{\mathbf{K}} \preceq \frac{3}{2} \mathbf{K}(l)$, and $\mathbf{B}(l)$ consists of $O(dR^2 \log^2 d)$ rows of $\mathbf{A}(l)$. Combining this with the error probability in the induction hypothesis shows that it holds for l as well.

For final output, combining $\gamma(l) \leq \lambda_{\min}$ with Lemma 4.2 gives that $\sqrt{\frac{1}{2}} \mathbf{B}(l)$ is a $2d$ -stretch base for $\mathbf{A}$. Applying Lemmas 2.8 and Lemma 2.1 once more to $\tilde{\tau}$ gives the bound on $\tilde{\mathbf{K}}$.

The last call to SAMPLE produces a $\mathbf{B}$ with $O(dR^2 \log^2 d / \epsilon^2)$ scaled rows of $\mathbf{A}$, which means it costs $O(d^\omega R^2 \log^2 d / \epsilon^2)$ to evaluate $\tilde{\mathbf{K}}$. It remains to bound the cost of approximating stretches and sampling from the $O(\log_R(\lambda_{upper} / \lambda_{lower}))$ iterations. As $\mathbf{B}(l)$ has $O(dR^2 \log^2 d)$ rows, Lemma 2.8 gives a total of $O((\mathbf{nnz}(\mathbf{A}) + \mathbf{nnz}(\mathbf{B}) + d^2) \log_R n + d^\omega R^2 \log^2 d)$ time. Combining this terms with the iteration count gives the overall running time. $\blacksquare$

In order to obtain a running time of $O(\mathbf{nnz}(\mathbf{A}) + d^{\omega+\alpha})$ for some constant $\alpha > 0$, we can set R to $d^{\alpha/2}$. Theorem 4.4 gives that ROWSAMPLE runs in $O((\mathbf{nnz}(\mathbf{A}) + d^2) \log_d n + d^{\omega+\alpha} \log^2 d \log_d(\lambda_{upper} / \lambda_{lower}) + d^{\omega+\alpha} \log^2 d \epsilon^{-2})$ time, and with probability at least $1 - \delta$ returns $\mathbf{B}$ with $O(d^{1+\alpha} \log^2 d \epsilon^{-2})$ rescaled rows of $\mathbf{A}$. When n and $\lambda_{upper} / \lambda_{lower}$ are both **poly**(d), the running time simplifies to $O(\mathbf{nnz}(\mathbf{A}) + d^{\omega+\alpha} \epsilon^{-2})$.

The pseudocode as given in Algorithm 3 needs values for λ_{upper} and λ_{lower} given as input. This requirement can be relaxed in several ways. Taking the Frobenius norm of $\mathbf{A}$, $\|\mathbf{A}\|_2^2$ gives a factor d approximation of $\lambda_{\max}$. Furthermore, if we know the rank of $\mathbf{A}$, we can check whether $\gamma(l) < \lambda_{\min}$ by computing the eigenvalues of $\mathbf{K}(l)$ and comparing against the smallest one.

5 Remarks

Aside from giving the state-of-art running times for approximating $\mathbf{A}^T \mathbf{A}$, each row in the output of our algorithms is a scaled copy of some row of $\mathbf{A}$. This means any specialized structure for rows of $\mathbf{A}$ are preserved in $\mathbf{B}$. The algorithm described in Section 4 has the further advantage that all the intermediate matrices also consists of only rows of $\mathbf{A}$ and a minor of the identity matrix. In practice, this means any uniform sparsity in $\mathbf{A}$ would also exist in these intermediate problems. In situations where all columns of $\mathbf{A}$ is sparse, iterative methods can solve the $d \times d$ linear systems in times faster than d^ω . This speedup is difficult to quantify without the introduction of additional parameters, and there are few general bounds for iterative methods with sparse systems. As a result, we believe this additional property is of more practical value and should be examined experimentally.

We believe that removing the factor of d^α , as well as extending these approaches computing low-rank approximations are natural directions for future work.

References

- [AB12] Haim Avron and Christos Boutsidis. Faster subset selection for matrices and applications. *CoRR*, abs/1201.0127, 2012. 2.2
- [ACD⁺10] Emmanuel Agullo, Camille Coti, Jack Dongarra, Thomas Herault, and Julien Langou. QR Factorization of Tall and Skinny Matrices in a Grid Computing Environment. In *24th IEEE International Parallel and Distributed Processing Symposium (IPDPS 2010)*, Atlanta, États-Unis, 2010. 1
- [Ach01] Dimitris Achlioptas. Database-friendly random projections. In *Proceedings of the twentieth ACM SIGMOD-SIGACT-SIGART symposium on Principles of database systems*, PODS '01, pages 274–281, New York, NY, USA, 2001. ACM. 2.3
- [AMT10] Haim Avron, Petar Maymounkov, and Sivan Toledo. Blendenpik: Supercharging lapack’s least-squares solver. *SIAM J. Scientific Computing*, pages 1217–1236, 2010. 1, 2.2
- [AT11] Haim Avron and Sivan Toledo. Effective stiffness: Generalizing effective resistance sampling to finite element matrices. *CoRR*, abs/cs/1110.4437, 2011. 2.1, 2.1
- [AW02] Rudolf Ahlswede and Andreas Winter. Strong converse for identification via quantum channels. *IEEE Transactions on Information Theory*, 48(3):569–579, 2002. 2.1, 2.1
- [Axe94] Owe Axelsson. *Iterative Solution Methods*. Cambridge University Press, New York, NY, 1994. 1
- [BHM00] William L. Briggs, Van Emden Henson, and Steve F. McCormick. *A multigrid tutorial: second edition*. Society for Industrial and Applied Mathematics, 2000. 3
- [CG11] Paul G. Constantine and David F. Gleich. Tall and skinny QR factorizations in mapreduce architectures. In *Proceedings of the second international workshop on MapReduce and its applications*, MapReduce '11, pages 43–50, New York, NY, USA, 2011. ACM. 1
- [CW12] Kenneth L. Clarkson and David P. Woodruff. Low rank approximation and regression in input sparsity time. *CoRR*, abs/0911.0547, 2012. 1, 2
- [DM10] Petros Drineas and Michael W. Mahoney. Effective resistances, statistical leverage, and applications to linear equation solving. 2010. 2.1
- [DMIMW12] Petros Drineas, Malik Magdon-Ismael, Michael W. Mahoney, and David P. Woodruff. Fast approximation of matrix coherence and statistical leverage. *ICML*, 2012. 1, 2.3, 2.3
- [DMM06] Petros Drineas, Michael W. Mahoney, and S. Muthukrishnan. Sampling algorithms for l2 regression and applications. In *Proceedings of the seventeenth annual ACM-SIAM symposium on Discrete algorithm*, SODA '06, pages 1127–1136, New York, NY, USA, 2006. ACM. 1
- [Har11] Nicholas Harvey. C&O 750: Randomized algorithms, winter 2011, lecture 11 notes. 2011. 2.1, 2.1
- [IM98] Piotr Indyk and Rajeev Motwani. Approximate nearest neighbors: towards removing the curse of dimensionality. In *STOC '98: Proceedings of the thirtieth annual ACM symposium on Theory of computing*, pages 604–613, New York, NY, USA, 1998. ACM. 2.3, 2.6
- [IW12] I. C. F. Ipsen and T. Wentworth. The Effect of Coherence on Sampling from Matrices with Orthonormal Columns, and Preconditioned Least Squares Problems. *CoRR*, abs/1203.4809, 2012. 2.1

-
- [KLP12] Ioannis Koutis, Alex Levin, and Richard Peng. Improved spectral sparsification and numerical algorithms for SDD matrices. In *29th International Symposium on Theoretical Aspects of Computer Science, STACS 2012*, pages 266–277, 2012. [1](#), [2.2](#), [2.2](#), [2.3](#)
 - [MI10] Malik Magdon-Ismael. Row sampling for matrix algorithms via a non-commutative bernstein bound. *CoRR*, abs/1008.0587, 2010. [1](#)
 - [RV07] Mark Rudelson and Roman Vershynin. Sampling from large matrices: An approach through geometric functional analysis. *J. ACM*, 54(4):21, 2007. [2.1](#), [2.1](#)
 - [Saa96] Yousef Saad. *Iterative Methods for Sparse Linear Systems*. <http://www-users.cs.umn.edu/~saad/books.html>, 1996. [1](#)
 - [SLHD10] Fengguang Song, Hatem Ltaief, Bilel Hadri, and Jack Dongarra. Scalable tile communication-avoiding QR factorization on multicore cluster systems. In *Proceedings of the 2010 ACM/IEEE International Conference for High Performance Computing, Networking, Storage and Analysis, SC '10*, pages 1–11, Washington, DC, USA, 2010. IEEE Computer Society. [1](#)
 - [SS08] Daniel A. Spielman and Nikhil Srivastava. Graph sparsification by effective resistances. In *Proceedings of the 40th Annual ACM Symposium on Theory of Computing (STOC)*, pages 563–568, 2008. [1](#), [2.1](#), [2.2](#), [2.2](#), [2.3](#), [2.3](#)
 - [Str93] Gilbert Strang. *Introduction to Linear Algebra*. Wellesley-Cambridge Press, 1993. [1](#)
 - [SW09] Daniel A. Spielman and Jaehoon Woo. A note on preconditioning by low-stretch spanning trees. *CoRR*, abs/0903.2816, 2009. [2.2](#)
 - [Ver09] R. Vershynin. A note on sums of independent random matrices after ahlswede-winter. 2009. [2.1](#)

A Proofs From Section 2

Proof of Fact 2.2:

$$\begin{aligned}
\sum_{i=1}^n \mathcal{STR}_{\mathbf{K}}(\mathbf{a}_i) &= \sum_{i=1}^n \mathbf{a}_i \mathbf{K}^+ \mathbf{a}_i^T \\
&= \sum_{i=1}^n \text{tr} [\mathbf{K}^+ \mathbf{a}_i^T \mathbf{a}_i] \\
&= \text{tr} \left[\mathbf{K}^+ \sum_{i=1}^n \mathbf{a}_i^T \mathbf{a}_i \right] \\
&= \text{tr} [\mathbf{K}^+ \mathbf{K}] = r
\end{aligned} \tag{1.21}$$

■

Proof of Fact 2.3: Note that both the generalized definition of leverage scores and the Forbenius norm works on each row independently. Therefore it suffices to prove this when $\mathbf{A}'$ has a single row, aka. $\mathbf{A}' = \mathbf{a}$. In this case the cyclic property of trace gives:

$$\begin{aligned}
\mathcal{STR}_{\mathbf{K}}(\mathbf{a}) &= \mathbf{a} \mathbf{K}^+ \mathbf{a}^T \\
&= \mathbf{a} \mathbf{K}^{+1/2} \mathbf{K}^{+1/2} \mathbf{a}^T \\
&= \|\mathbf{K}^{+1/2} \mathbf{a}^T\|_2^2
\end{aligned} \tag{1.22}$$

■

Proof of Lemma 2.5: The condition given implies that the null spaces of $\mathbf{K}$ and $\tilde{\mathbf{K}}$ are identical, giving:

$$\mathbf{K}^+ \preceq \tilde{\mathbf{K}}^+ \preceq \kappa \mathbf{K}^+ \tag{1.23}$$

Applying this to the vector $\mathbf{x}$ gives:

$$\mathbf{x}\mathbf{K}^+\mathbf{x}^T \leq \mathbf{x}\tilde{\mathbf{K}}^+\mathbf{x}^T \leq \kappa\mathbf{x}\tilde{\mathbf{K}}^+\mathbf{x}^T \quad (1.24)$$

The first inequality gives Part 1, while summing the second inequality over all rows gives Part 2. $\blacksquare$

B Proofs About Projections and Prolongations

Proof of Lemma 3.1: Let $\mathbf{K} = \mathbf{A}^T \mathbf{A}$, by Fact 2.3, we have that

$$\mathcal{STR}_{\mathbf{A}}(\mathbf{A}_{(b)}) = \|\mathbf{A}_{(b)}\mathbf{K}^{+1/2}\|_F^2$$

and

$$\mathcal{STR}_{\mathbf{A}}(\mathbf{A}(\text{short})_{(b)}) = \|\mathbf{A}(\text{short})_{(b)}\mathbf{K}^{+1/2}\|_F^2$$

Furthermore, since $\mathbf{A}(\text{short})_{(b)} = \mathbf{U}_{(b)}\mathbf{A}_{(b)}$, we have:

$$\begin{aligned} \mathcal{STR}_{\mathbf{A}}(\mathbf{A}(\text{short})_{(b)}) &= \|\mathbf{A}(\text{short})_{(b)}\mathbf{K}^{+1/2}\|_F^2 \\ &= \|\mathbf{U}_{(b)}\mathbf{A}_{(b)}\mathbf{K}^{+1/2}\|_F^2 \end{aligned} \quad (2.25)$$

Let $\mathbf{Y}$ denote $\mathbf{A}_{(b)}\mathbf{K}^{+1/2}$, then we have:

$$\begin{aligned} &\Pr\left(\mathcal{STR}_{\mathbf{A}}(\mathbf{A}(\text{short})_{(b)}) < \frac{k}{R}\mathcal{STR}_{\mathbf{A}}(\mathbf{A}_{(b)})\right) \\ &= \Pr\left(\|\mathbf{U}_{(b)}\mathbf{Y}\|_F^2 < \frac{k}{R}\|\mathbf{Y}\|_F^2\right) \end{aligned} \quad (2.26)$$

Consider each column of $\mathbf{Y}$, $\mathbf{y}_{:,j}$. By Lemma 2.6, we have that:

$$\begin{aligned} &\Pr\left(\|\mathbf{U}_{(b)}\mathbf{y}_{:,j}\|_2^2 < \frac{k}{\beta}\|\mathbf{y}_{:,j}\|_2^2\right) \\ &\leq \exp\left(-\frac{k}{2}(\beta - (1 + \ln \beta))\right) \end{aligned} \quad (2.27)$$

Setting $\beta = R$, this value can be bounded by $O(k)\exp(-\frac{k}{2}(\beta^{-1} - 1 - \ln \beta)) \leq O(k)R^{-k/4}$. As $R \geq d^\alpha$, setting $k = 4(c+1)/\alpha$ gives that this probability is at most d^{-c-1} . Applying union bound over all d columns gives that with probability $1 - d^{-c}$ we have that for all columns, $\|\mathbf{U}_{(b)}\mathbf{y}_{:,j}\|_2^2 \geq \frac{k}{\beta}\|\mathbf{y}_{:,j}\|_2^2$. The overall bound then follows from fact that $\|\mathbf{Y}\|_F^2 = \sum_{j=1}^d \|\mathbf{y}_{:,j}\|_2^2$. $\blacksquare$

Proof of Lemma 3.2: By definition of spectral norm, we have $\|\mathbf{U}_{(b)}^T \mathbf{U}_{(b)}\| \leq \|\mathbf{U}_{(b)}\|_F^2$, and in turn:

$$\|\mathbf{U}_{(b)}\mathbf{y}\|_2^2 \leq \|\mathbf{U}_{(b)}\|_F^2 \cdot \|\mathbf{y}\|_2^2 \quad (2.28)$$

for any vector $\mathbf{y}$.

Let $\mathbf{x}$ be any vector in $\mathbb{R}^d$. Substituting in $\mathbf{A}(\text{short})_{(b)} = \mathbf{U}_{(b)}\mathbf{A}_{(b)}$ gives:

$$\begin{aligned}
& \mathbf{x}^T \mathbf{A}(\text{short})_{(b)}^T \mathbf{A}(\text{short})_{(b)} \mathbf{x} \\
&= \|\mathbf{A}(\text{short})_{(b)} \mathbf{x}\|_2^2 \\
&\leq \|\mathbf{U}_{(b)}\|_F^2 \cdot \|\mathbf{A}_{(b)} \mathbf{x}\|_2^2 \quad \text{By Equation 2.28} \\
&= \|\mathbf{U}_{(b)}\|_F^2 \cdot \mathbf{x}^T \mathbf{A}_{(b)}^T \mathbf{A}_{(b)} \mathbf{x}
\end{aligned} \tag{2.29}$$

■

Proof of Lemma 3.3: Consider an orthonormal basis for the range space of $\mathbf{C}$, $\mathbf{v}_1 \dots \mathbf{v}_{\text{rank}(\mathbf{C})}$. Since $\mathbf{C} + \mathbf{D} \succeq \mathbf{C}$, this basis can be extended to an orthonormal basis to the range space of $\mathbf{C} + \mathbf{D}$ by adding $\mathbf{v}_{\text{rank}(\mathbf{C})+1} \dots \mathbf{v}_{\text{rank}(\mathbf{C}+\mathbf{D})}$. It suffices to prove the claim under this basis system. Here $\mathbf{C}$ and $\mathbf{D}$ can be rewritten as:

$$\mathbf{C} = \begin{bmatrix} \mathbf{C}_{11} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \quad \mathbf{D} = \begin{bmatrix} \mathbf{D}_{11} & \mathbf{D}_{12} \\ \mathbf{D}_{12}^T & \mathbf{D}_{22} \end{bmatrix} \tag{2.30}$$

Where $\mathbf{C}_{11}$ and $\mathbf{D}_{22}$ are strictly positive definite. Furthermore, since $\mathbf{D}$ is positive semi-definite we have that $\mathbf{D}_{11} - \mathbf{D}_{12}\mathbf{D}_{22}^{-1}\mathbf{D}_{12}$ is also positive semi-definite. For any vector $\mathbf{x}$, $\Pi\mathbf{x}$ gives a vector that's non-zero only in the first $\text{rank}(\mathbf{C})$ entries. Let this part be $\mathbf{x}_1$. Then evaluating $(\mathbf{C} + \mathbf{D})^+ \Pi\mathbf{x} = [\mathbf{y}_1; \mathbf{y}_2]$ becomes solving the following system:

$$(\mathbf{C}_{11} + \mathbf{D}_{11})\mathbf{y}_1 + \mathbf{D}_{12}\mathbf{y}_2 = \mathbf{x}_1 \tag{2.31}$$

$$\mathbf{D}_{12}^T \mathbf{y}_1 + \mathbf{D}_{22}\mathbf{y}_2 = \mathbf{0} \tag{2.32}$$

The second equation gives $\mathbf{y}_2 = -\mathbf{D}_{22}^{-1}\mathbf{D}_{12}^T \mathbf{y}_1$. Substituting it into the first one gives:

$$(\mathbf{C}_{11} + \mathbf{D}_{11} - \mathbf{D}_{12}\mathbf{D}_{22}^{-1}\mathbf{D}_{12}^T)\mathbf{y}_1 = \mathbf{x}_1 \tag{2.33}$$

Note that this is the same as taking the partial Cholesky factorization onto the range space of Π . Combining things gives:

$$\mathbf{x}^T \Pi(\mathbf{C} + \mathbf{D})^+ \Pi\mathbf{x} = \mathbf{x}_1^T (\mathbf{C}_{11} + \mathbf{D}_{11} - \mathbf{D}_{12}\mathbf{D}_{22}^{-1}\mathbf{D}_{12}^T)^{-1} \mathbf{x}_1 \tag{2.34}$$

Since $\mathbf{D}_{11} - \mathbf{D}_{12}\mathbf{D}_{22}^{-1}\mathbf{D}_{12}^T$ is positive definite, we have $\mathbf{C}_{11} \preceq \mathbf{C} + \mathbf{D}_{11} - \mathbf{D}_{12}\mathbf{D}_{22}^{-1}\mathbf{D}_{12}^T$ and therefore:

$$\begin{aligned}
\mathbf{x}^T \Pi(\mathbf{C} + \mathbf{D})^+ \Pi\mathbf{x} &= \mathbf{x}_1^T (\mathbf{C}_{11} + \mathbf{D}_{11} - \mathbf{D}_{12}\mathbf{D}_{22}^{-1}\mathbf{D}_{12}^T)^{-1} \mathbf{x}_1^T \\
&\leq \mathbf{x}_1^T \mathbf{C}_{11}^{-1} \mathbf{x}_1^T \\
&= \mathbf{x}^T \Pi\mathbf{C}^+ \Pi\mathbf{x}
\end{aligned} \tag{2.35}$$

■